

Limited dispersal in mobile hunter–gatherer Baka Pygmies

Paul Verdu, Raphaël Leblois, Alain Froment, Sylvain Théry, Serge Bahuchet, François Rousset, Evelyne Heyer and Renaud Vitalis

Biol. Lett. published online 28 April 2010
doi: 10.1098/rsbl.2010.0192

References

[This article cites 14 articles, 1 of which can be accessed free](#)

<http://rsbl.royalsocietypublishing.org/content/early/2010/04/23/rsbl.2010.0192.full.html#ref-list-1>

P<P

Published online 28 April 2010 in advance of the print journal.

Subject collections

Articles on similar topics can be found in the following collections

[ecology](#) (1381 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To subscribe to *Biol. Lett.* go to: <http://rsbl.royalsocietypublishing.org/subscriptions>

Limited dispersal in mobile hunter–gatherer Baka Pygmies

Paul Verdu^{1,2,*}, Raphaël Leblois³, Alain Froment⁴, Sylvain Théry², Serge Bahuchet², François Rousset⁵, Evelyne Heyer² and Renaud Vitalis^{2,†}

¹Department of Human Genetics, University of Michigan, Ann Arbor, MI, USA

²MNHN–CNRS, Université Paris 7, UMR 7206 ‘Ecoanthropology and Ethnobiology’, Paris, France

³MNHN–CNRS, UMR 7205 ‘Origine, Structure et Evolution de la Biodiversité’, Paris, France

⁴IRD–MNHN, UMR 208 ‘Patrimoines locaux’, Paris, France

⁵Université Montpellier 2–CNRS, UMR 5554 ‘Institut des Sciences de l’Evolution’, Montpellier, France

*Author for correspondence (verdu@umich.edu).

†Present address: CNRS–INRA, UMR CBGP (INRA–IRD–CIRAD–Montpellier SupAgro), Campus International de Baillarguet, Montpellier-sur-Lez, France.

Hunter–gatherer Pygmies from Central Africa are described as being extremely mobile. Using neutral genetic markers and population genetics theory, we explored the dispersal behaviour of the Baka Pygmies from Cameroon, one of the largest Pygmy populations in Central Africa. We found a strong correlation between genetic and geographical distances: a pattern of isolation by distance arising from limited parent–offspring dispersal. Our study suggests that mobile hunter–gatherers do not necessarily disperse over wide geographical areas.

Keywords: African Pygmies; dispersal; isolation by distance; microsatellites; population genetics

1. INTRODUCTION

The correlation between genetics and geography has been extensively studied at a worldwide scale to better characterize the distribution of human genetic diversity and estimate key parameters of modern humans’ expansion worldwide (e.g. Ramachandran *et al.* 2005; Liu *et al.* 2006; DeGiorgio *et al.* 2009). However, none of these studies have characterized parent–offspring dispersal and therefore, the dispersal mechanisms underlying the geographical distribution of human genetic diversity remain unclear.

Ethnologists have shown the great variability of mobility across human populations with different lifestyles and modes of subsistence. For example, foragers have been shown to be much more mobile than farmers (MacDonald & Hewlett 1999). However, the relationship between mobility and effective dispersal (characterized by the distances between children’s and parents’ birthplaces) often remains unknown. To explore this aspect, we studied African Pygmies, which represent the largest hunter–gatherer group of populations worldwide. African Pygmies are described as being very mobile within the rainforest for seasonal mobility and socio-economic activities (Bahuchet 1992). Demographic data in

an Aka Pygmy population further showed large mating ranges and large distances between birthplaces and places of residence (Cavalli-Sforza & Hewlett 1982). This suggests that such mobile behaviour should translate into long distances between children’s and parents’ birthplaces, but this assumption has never been explicitly tested owing to limited demographic data (Cavalli-Sforza 1986). In this context, population genetics can provide indirect estimates of effective dispersal in these hunter–gatherers. Indeed, isolation by distance theory predicts that at equilibrium between drift, migration and mutation, the regression of F -statistics estimates on the logarithm of geographical distances provides a robust estimate of the neighbourhood size, which is proportional to the population density and the second moment of parent–offspring distance, if individuals are sampled at a short geographical scale (Rousset 1997).

Here, we report estimates of effective dispersal based on the relationship between genetic and geographical distances in three groups of the Baka Pygmies from Cameroon. This is, to the best of our knowledge, the only genetic data for such a small geographical scale in African Pygmy populations, which provides an opportunity to infer effective dispersal using the genetics of these mobile hunter–gatherers.

2. MATERIAL AND METHODS

The Baka from Cameroon represent one of the largest Pygmy populations with 35 000 individuals occupying 75 000 km² in the rainforest (figure 1). These values, which were compiled from ethnographic data (Vallois & Marquer 1976; Dhellemmes & Macaigne 1985; Cavalli-Sforza 1986; Sato 1992; Abéga 1998; Tsuru 2001), provide an estimate of the population density of $D = 0.47$ individuals · km⁻².

We consider 87 Baka adults genotyped at 28 independent tetra-nucleotide microsatellite loci (Verdu *et al.* 2009). We visited Baka settlements along three transects of approximately 50 km each (figure 1), asking volunteers to gather for DNA sampling at a single location for each transect. Frequent movements between places of temporary residence, as well as hospitality rules among Pygmies that assimilate visitors as residents during their visit, make it difficult to determine whether an individual met in a village is a resident or a visitor (Cavalli-Sforza 1986; Bahuchet 1992). Therefore, the sampling location, or the location where individuals were first met, are probably a poor predictor of Pygmies’ location after dispersal and these data were discarded. Instead, we considered that birthplaces provided more robust data to study Pygmies’ dispersal. After collecting each individual’s birthplace, we went back on the road to determine the geographical coordinates of these locations. Each donor provided appropriate informed oral consent.

Geographically limited dispersal across generations in two-dimensional habitats results in a linear relationship between pairwise genetic distances a_{ij} , an analogue to $F_{ST}/(1 - F_{ST})$ calculated between pairs of individuals (Rousset 2000), and the logarithm of geographical distance. We used the software package GENEPOP’007 (Rousset 2008) to calculate the slope of this linear regression, which, at a small geographic scale, provides a robust estimator of $1/(4D\sigma^2)$, where D is the effective population density and σ^2 is the second moment of parent–offspring axial distance (Rousset 1997). We tested the significance of the correlation using a Mantel test (Mantel 1967) with 30 000 permutations. First, we considered all pairs of individuals from the three geographically distant Baka groups (figure 1). Second, we considered only pairs of individuals born within the same group and discarded pairs of individuals born in different groups. To that end, we wrote a R script (R Development Core Team 2007), available upon request, which modified the Mantel test to calculate rank correlation coefficients and to permute the pairwise distances within groups only.

3. RESULTS AND DISCUSSION

Considering all possible pairs of Baka individuals, we found a significant positive correlation between pairwise genetic distance and the logarithm of geographical distance ($p = 0.010$): individuals born




Figure 1. Geographical distribution of the 87 Baka Pygmies sampled in southeastern Cameroon. In each Baka group, birthplaces of sampled individuals are shown with sample sizes given in parentheses. Other Pygmy peopling areas inferred from our ethnographical fieldworks are shown. Map sources: Global Land Cover Facility.

nearby are more genetically related than individuals born further away from each other. The slope of regression between the genetic and geographical distances equalled 0.0027 (95% CI 0.0007–0.0046; see figure 2). We estimated from this slope that $4\pi D\sigma^2 = 373$ individuals and, using $D = 0.47$ individuals \cdot km $^{-2}$, we obtained $\sigma^2 = 63.2$ km 2 . But theory shows that mutation wipes out the linear increase in genetic differentiation with geographical distance, if the distances between samples are larger than $d_{\max} = 0.56 \sigma / \sqrt{2\mu}$, where μ is the mutation rate (Rousset 1997, 2004). Here, the maximum distance between any two individuals' birthplace was 296 km, a value much larger than $d_{\max} = 120$ km, assuming $\mu = 7 \times 10^{-4}$ (Zhivotovsky *et al.* 2003). It is therefore likely that considering the full range of geographical distances in this first analysis leads to an overestimate of σ^2 .

Therefore, in the second analysis, we discarded all pairs of individuals born in different Baka groups and considered only pairs of individuals born nearby. We then found a stronger increase of genetic differentiation with geographical distance (slope = 0.0137; 95% CI 0.0038–0.0265; $p = 0.004$; figure 2) providing an estimate of $4D\pi\sigma^2 = 73$ individuals, which gave $\sigma^2 = 12.4$ km 2 . Here, the maximum distance between two sampled individuals' birthplaces (26 km) was shorter than $d_{\max} = 53$ km, indicating that this estimate of σ^2 should no longer be biased by mutation. But we cannot exclude that reducing




Figure 2. Correlation between genetic differentiation and the logarithm of geographical distances among Baka Pygmies. Multilocus estimates of pairwise differentiation (\hat{a}_r) are plotted against the logarithm of geographical distances (in kilometres). The linear regression considering all pairs of individuals is $y = 0.0027x - 0.0153$ (in blue). The linear regression considering only pairs of individuals born within the same group is $y = 0.0137x - 0.1138$ (in red).

the sampling scale potentially excluded some long-distance migrants, which would lead to an underestimate of σ^2 (Rousset 2004).

Altogether, this suggests that the effective parent–offspring dispersal in the Baka Pygmies lies between 12.4 and 63.2 km 2 . These indirect estimates of effective dispersal average over male and female genetic contributions, since they are based on autosomal

data. Assuming that parent–offspring dispersal distances are exponentially distributed (Cavalli-Sforza & Hewlett 1982), then half of the offspring disperse at distances shorter than $\sigma \ln(2)/\sqrt{2} \approx 0.49 \sigma$. In this case, half of the Baka offspring disperse at a maximal distance between 1.7 and 3.9 km. However, using other distributions than the exponential for the dispersal distances would provide different interpretations. Therefore, dispersal distances estimated from σ^2 should be preferred here since they are independent of the (unknown) shape of the dispersal distribution.

Without quantitative demographic data on the Baka's mobility, we used the available demographic data from the Aka Pygmies from the Central African Republic (Cavalli-Sforza & Hewlett 1982) for comparison. We calculated the second moment of the distribution of distances between birthplaces and places of residence, an estimate of σ^2 from demographic data. From table 5 in Cavalli-Sforza & Hewlett (1982), we computed σ^2 as $\sum_i i^2 p_i$, where p_i is the proportion of individuals whose birthplace and place of residence are separated by i kilometres. We found $\sigma^2 = 3599 \text{ km}^2$ for men and 4061 km^2 for women (3683 km^2 on average). Hence, our estimates of the Baka's dispersal from genetic data are, respectively, 58.3 and 297 times lower than the average estimate of dispersal from demographic data in the Aka. How can this discrepancy be reconciled?

First, ethnologists show that mobility significantly differs across Pygmy populations from Central Africa (Bahuchet 1992). For instance, the mobility of the Bongo from Gabon probably decreased recently as opposed to the extended mobility of the Aka, since the Bongo nowadays live in permanent houses and widely practice agriculture (P. Verdu 2006–2007, unpublished results). In this context, the difference between our indirect estimates of dispersal in the Baka and the direct estimates in the Aka could result from differences in dispersal behaviour between these groups. Second, Cavalli-Sforza & Hewlett (1982) provide the distribution of distances between birthplaces and places of residence. This demographic data primarily reflects exploration behaviour rather than effective parent–offspring dispersal. To consistently compare demographic and genetic estimates, demographic estimates require an accurate knowledge of parents' and offspring's birthplaces, which is particularly challenging in Pygmies (Cavalli-Sforza 1986).

In conclusion, we found a strong signal of isolation by distance among the Baka Pygmies, a pattern owing to limited parent–offspring dispersal. Although our results do not challenge the view that hunter–gatherer Pygmies have frequent movements in their socio-economic area, we demonstrate that extended individual mobility does not necessarily reflect extended dispersal across generations. Limited effective dispersal may have reinforced genetic isolation among Pygmy populations, which could be a key mechanism explaining the strong genetic differentiation found among Western Central African Pygmies, despite their recent divergence from a single ancestral population about 2800 years ago (Verdu *et al.* 2009).

More generally, our findings also challenge the view that mobile hunter–gatherers disperse more than sedentary farmers. Hunter–gatherers are generally

described as much more mobile than agricultural populations, based on mating distances (see fig. 3 in MacDonald & Hewlett (1999)). Here, we show that Baka hunter–gatherers' dispersal is strongly localized, as is the effective dispersal previously estimated in the horticulturalist Gainj- and Kalam-speakers from New Guinea ($\sigma^2 = 1.41 \text{ km}^2$; Rousset 1997), the only other human genetic data collected at the appropriate geographical scale for applying Rousset's (1997) regression method (Long *et al.* 1987). This suggests that, despite their very mobile behaviour, foragers do not necessarily disperse more than farmers.

Research and sampling authorizations were obtained from the ethical committees of both the French and the Cameroonian governments.

The authors warmly thank all donors from Cameroon. We thank Frédéric Austerlitz, Michael DeGiorgio, Ethan Jewett, Trevor Pemberton, Noah Rosenberg and two anonymous reviewers for useful suggestions. This work was funded by the CNRS, the ACI Prosodie and the ANR (grant 05-BLAN-0400-01 MPRAs; programme BLANC 'EMILE' NT09-611697).

- Abéga, S. C. 1998 *Pygmées Baka: le droit à la différence*. Yaoundé, Cameroon: INADES-formation Cameroun.
- Bahuchet, S. 1992 Spatial mobility and access to the resources among the African Pygmies. In *Mobility and territoriality* (eds M. J. Casimir & A. Rao), pp. 205–257. New York, NY; Oxford, UK: Berg.
- Cavalli-Sforza, L. L. 1986 *African pygmies*. Orlando, FL: Academic Press.
- Cavalli-Sforza, L. L. & Hewlett, B. 1982 Exploration and mating range in African Pygmies. *Ann. Hum. Genet.* **46**, 257–270. (doi:10.1111/j.1469-1809.1982.tb00717.x)
- DeGiorgio, M., Jakobsson, M. & Rosenberg, N. A. 2009 Explaining worldwide patterns of human genetic variation using a coalescent-based serial founder model of migration outward from Africa. *Proc. Natl Acad. Sci. USA* **106**, 16 057–16 062. (doi:10.1073/pnas.0903341106)
- Dhellemmes, I. & Macaigne, P. 1985 *Le père des pygmées*. Aventure vécue. Paris, France: Flammarion.
- Liu, H., Prugnolle, F., Manica, A. & Balloux, F. 2006 A geographically explicit genetic model of worldwide human-settlement history. *Am. J. Hum. Genet.* **79**, 230–237. (doi:10.1086/505436)
- Long, J. C., Smouse, P. E. & Wood, J. W. 1987 The allelic correlation structure of Gainj- and Kalam-speaking people. II. The genetic distance between population subdivisions. *Genetics* **117**, 273–283.
- MacDonald, D. H. & Hewlett, B. S. 1999 Reproductive interests and forager mobility. *Curr. Anthropol.* **40**, 501–523. (doi:10.1086/200047)
- Mantel, N. 1967 The detection of disease clustering and a generalized regression approach. *Cancer Res.* **27**, 209–220.
- R Development Core Team 2007 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Ramachandran, S., Deshpande, O., Roseman, C. C., Rosenberg, N. A., Feldman, M. W. & Cavalli-Sforza, L. L. 2005 Support from the relationship of genetic and geographic distance in human populations for a serial founder effect originating in Africa. *Proc. Natl Acad. Sci. USA* **102**, 15942–15947. (doi:10.1073/pnas.0507611102)

- Rousset, F. 1997 Genetic differentiation and estimation of gene flow from F -statistics under isolation by distance. *Genetics* **145**, 1219–1228.
- Rousset, F. 2000 Genetic differentiation between individuals. *J. Evol. Biol.* **13**, 58–62. (doi:10.1046/j.1420-9101.2000.00137.x)
- Rousset, F. 2004 *Genetic structure and selection in subdivided populations*. Monographs in population biology. Princeton, NJ: Princeton University Press.
- Rousset, F. 2008 GENEPOP '007: a complete re-implementation of the GENEPOP software for Windows and Linux. *Mol. Ecol. Resour.* **8**, 103–106. (doi:10.1111/j.1471-8286.2007.01931.x)
- Sato, H. 1992 Notes on the distribution and settlement pattern of hunter–gatherers in Northwestern Congo. *African Study Monographs* **13**, 203–216.
- Tsuru, D. 2001 Generation and transaction processes in the spirit ritual of the Baka Pygmies in Southeast Cameroon. *Afric. Stud. Monographs, Suppl.* **27**, 103–123.
- Vallois, H. V. & Marquer, P. 1976 *Les Pygmées Baka du Cameroun: anthropologie et ethnographie. Mémoires du Mus. Nat. Hist. Nat., Série A, Tome C*. Paris, France: Muséum National d'Histoire Naturelle.
- Verdu, P. *et al.* 2009 Origins and genetic diversity of pygmy hunter-gatherers from Western Central Africa. *Curr. Biol.* **19**, 312–318. (doi:10.1016/j.cub.2008.12.049)
- Zhivotovsky, L. A., Rosenberg, N. A. & Feldman, M. W. 2003 Features of evolution and expansion of modern humans, inferred from genomewide microsatellite markers. *Am. J. Hum. Genet.* **72**, 1171–1186. (doi:10.1086/375120)